


Catalytic Reactions of 1,1-Diborylalkanes and (Diborylmethyl)metallic Species

Seung Hwan Cho

Department of Chemistry, POSTECH, E-mail: seunghwan@postech.ac.kr

Organoborons are highly versatile intermediates in organic chemistry because they can serve as versatile platforms to afford various synthetically important chiral building blocks via stereospecific carbon-carbon and carbon-heteroatom bond forming reactions. Over the past few years, 1,1-diborylalkanes, which contain two identical boron groups at the same sp^3 carbon center, have been emerged as new types of synthetic precursors. Such reagents provide significant advantages for increasing molecular diversity and complexity through stereospecific and/or iterative carbon-carbon and carbon-heteroatom bond-forming reactions.

In this seminar, the details about our recent findings for the catalytic enantioselective transformations of 1,1-diborylalkanes and (diborylmethyl)metallic species will be presented.¹⁻⁵



References

1. J. Kim, S. Park, J. Park and **S. H. Cho*** *Angew. Chem. Int. Ed.* **2016**, *55*, 1498.
2. J. Kim, K. Ko and **S. H. Cho*** *Angew. Chem. Int. Ed.* **2017**, *56*, 11584.
3. J. Kim, M. Shin and **S. H. Cho*** *ACS Catal.* **2019**, *9*, 850.
4. Y. Lee, J. Park and **S. H. Cho*** *Angew. Chem. Int. Ed.* **2018**, *57*, 12930.
5. J. Kim and **S. H. Cho*** *ACS Catal.* **2019**, *9*, 230.